

**APPROXIMATION OF SOLUTIONS OF THE FORCED
DUFFING EQUATION WITH NONLOCAL DISCONTINUOUS
TYPE
INTEGRAL BOUNDARY CONDITIONS**

AHMED ALSAEDI

ABSTRACT. A generalized quasilinearization technique is applied to obtain a sequence of approximate solutions converging monotonically and quadratically to the unique solution of the forced Duffing equation with nonlocal discontinuous type integral boundary conditions.

1. INTRODUCTION

Integral boundary conditions for evolution problems have various applications in chemical engineering, thermoelasticity, underground water flow and population dynamics, see for example [16, 17, 24]. In fact, boundary value problems involving integral boundary conditions have received considerable attention, see for instance, [3, 10], [12]–[15], [18, 19, 26] and the references therein. In a recent reference [2], Ahmad, et. al. discussed the existence and uniqueness of the solutions of a boundary value problem with discontinuous type integral boundary conditions.

The monotone iterative technique coupled with the method of upper and lower solutions [5, 8, 20, 23, 25] manifests itself as an effective and flexible mechanism that offers theoretical as well as constructive existence results in a closed set, generated by the lower and upper solutions. In general, the convergence of the sequence of approximate solutions given by the monotone iterative technique is at most linear [11, 21]. To obtain a sequence of approximate solutions converging quadratically, we use the method of quasilinearization (QSL) [9]. This method has been developed for a variety of problems [1, 4, 6, 7, 22]. In view of its diverse applications, this approach is quite an elegant and easier for application algorithms. To the best of our knowledge, the method of quasilinearization has not been developed for Duffing equation with nonlocal discontinuous type integral boundary conditions.

In this paper, we apply a quasilinearization technique to obtain the analytic approximation of the solution of the forced Duffing equation with nonlocal discontinuous type integral boundary conditions. In fact, we obtain a sequence of

2000 *Mathematics Subject Classification*: primary 34B10; secondary 34B15.

Key words and phrases: duffing equation, integral boundary conditions, quasilinearization, quadratic convergence.

Received December 12, 2007. Editor O. Došlý.

approximate solutions converging monotonically and quadratically to the unique solution of the problem at hand. The concept of nonlocal discontinuous integral boundary conditions corresponds to a situation when some forcing term is present at an arbitrary intermediate point of the boundary segment and thereby generates a discontinuity in the integral boundary conditions.

2. PRELIMINARIES

We consider the following boundary value problem

$$\begin{aligned}
 (2.1) \quad & u''(t) + \sigma u'(t) + f(t, u) = 0, \quad t \in [0, 1], \sigma \in \mathbb{R} \setminus \{0\}, \\
 & u(0) - \mu_1 u'(0) = g_1(u(\gamma)) + \int_0^{\gamma^-} q_1(u(s)) ds + \int_{\gamma^+}^1 q_1(u(s)) ds, \\
 & u(1) + \mu_2 u'(1) = g_2(u(\gamma)) + \int_0^{\gamma^-} q_2(u(s)) ds + \int_{\gamma^+}^1 q_2(u(s)) ds, \\
 & 0 < \gamma < 1,
 \end{aligned}$$

where $f: [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$, $g_i: \mathbb{R} \rightarrow \mathbb{R}$ ($i = 1, 2$) are continuous functions, q_i are continuous functions on $(0, \gamma)$ and $(\gamma, 1)$ and μ_i are nonnegative constants. The quasilinearization technique is applied to obtain a sequence of approximate solutions converging monotonically and quadratically to the unique solution of the problem (2.1).

Definition 2.1. A function $\alpha \in C^2[0, 1]$ is a lower solution of (2.1) if

$$\begin{aligned}
 & \alpha''(t) + \sigma \alpha'(t) + f(t, \alpha(t)) \geq 0, \quad t \in [0, 1], \\
 & \alpha(0) - \mu_1 \alpha'(0) \leq g_1(\alpha(\gamma)) + \int_0^{\gamma^-} q_1(\alpha(s)) ds + \int_{\gamma^+}^1 q_1(\alpha(s)) ds, \\
 & \alpha(1) + \mu_2 \alpha'(1) \leq g_2(\alpha(\gamma)) + \int_0^{\gamma^-} q_2(\alpha(s)) ds + \int_{\gamma^+}^1 q_2(\alpha(s)) ds.
 \end{aligned}$$

Similarly, $\beta \in C^2[0, 1]$ is an upper solution of (2.1) if the inequalities in the definition of lower solution are reversed.

Since the associated homogeneous problem of (2.1) has only the trivial solution, therefore, by Green's function method, the solution $u(t)$ of (2.1) can be written as

$$\begin{aligned}
 u(t) = & \frac{1}{(1 + \sigma\mu_1) - (1 - \sigma\mu_2)e^{-\sigma}} \left[((-1 + \sigma\mu_2)e^{-\sigma} + e^{-\sigma t}) \right. \\
 & \times \left\{ g_1(u(\gamma)) + \int_0^{\gamma^-} q_1(u(s)) ds + \int_{\gamma^+}^1 q_1(u(s)) ds \right\} \\
 & + ((1 + \sigma\mu_1) - e^{-\sigma t}) \left\{ g_2(u(\gamma)) + \int_0^{\gamma^-} q_2(u(s)) ds + \int_{\gamma^+}^1 q_2(u(s)) ds \right\} \Big] \\
 & + \int_0^1 G(t, s) f(s, u(s)) ds,
 \end{aligned}$$

where

$$G(t, s) = \Lambda \begin{cases} [(1 - \sigma\mu_2) - e^{\sigma(1-s)}][(1 + \sigma\mu_1) - e^{-\sigma t}], & 0 \leq t \leq s, \\ [(1 - \sigma\mu_2) - e^{\sigma(1-t)}][(1 + \sigma\mu_1) - e^{-\sigma s}], & s \leq t \leq 1, \end{cases}$$

$$\Lambda = \frac{e^{\sigma s}}{\sigma[(1 - \sigma\mu_2) - (1 + \sigma\mu_1)e^{\sigma}]}$$

Observe that $G(t, s) > 0$ on $(0, 1) \times (0, 1)$. We state the following results which lay a foundation to establish the main result. We omit the proof as the method of proof is similar to the one employed in [2].

Theorem 2.1. *Let α and β be lower and upper solutions of the boundary value problem (2.1) respectively. Let $f: [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ be such that $f_u(t, u) < 0$ and q_i are continuous functions on $(0, \gamma)$ and $(\gamma, 1)$ satisfying one sided Lipschitz condition: $q_i(u) - q_i(v) \leq L_i(u - v)$, $0 \leq L_i < 1$, $i = 1, 2$, and $g_i: \mathbb{R} \rightarrow \mathbb{R}$ are continuous functions satisfying one sided Lipschitz condition: $g_i(u) - g_i(v) \leq L_i^*(u - v)$, $0 \leq L_i^* < 1$, $i = 1, 2$. Then $\alpha(t) \leq \beta(t)$.*

Theorem 2.2. *Assume that α and β are lower and upper solutions of the boundary value problem (2.1) respectively such that $\alpha(t) \leq \beta(t)$. If $f: [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$, $g_i: \mathbb{R} \rightarrow \mathbb{R}$ are continuous functions and q_i are continuous functions on $(0, \gamma)$ and $(\gamma, 1)$ with g_i and q_i satisfying one sided Lipschitz condition, then there exists a solution $u(t)$ of (2.1) such that $\alpha(t) \leq u(t) \leq \beta(t)$, $t \in [0, 1]$.*

3. MAIN RESULT

Theorem 3.1. *Assume that*

- (A₁) α and $\beta \in C^2[0, 1]$ are respectively lower and upper solutions of (2.1) such that $\alpha(t) \leq \beta(t)$;
- (A₂) $f(t, u) \in C^2([0, 1] \times \mathbb{R})$ be such that $f_u(t, u) < 0$ and $(f_{uu}(t, u) + \phi_{uu}(t, u)) \geq 0$, where $\phi_{uu}(t, u) \geq 0$ for some continuous function $\phi(t, u)$ on $[0, 1] \times \mathbb{R}$;
- (A₃) q_i are continuous functions on $(0, \gamma)$ and $(\gamma, 1)$ satisfying $0 \leq q'_i(u) < 1$, and $(q''_i(u) + \chi''_i(u)) \geq 0$ with $\chi''_i \geq 0$, $i = 1, 2$;
- (A₄) $g_i \in C^2(\mathbb{R})$ be such that $0 \leq g'_i(u) < 1$ and $(g''_i(u) + \psi''_i(u)) \leq 0$ with $\psi''_i \leq 0$, $i = 1, 2$.

Then, there exists a sequence $\{\alpha_n\}$ of approximate solutions converging monotonically and quadratically to the unique solution of the problem (2.1).

Proof. Let $F: [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ and $G_i, K_i: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $F(t, u) = f(t, u) + \phi(t, u)$, $G_i(u) = g_i(u) + \psi_i(u)$, $K_i(u) = q_i(u) + \chi_i(u)$ so that $F_{uu}(t, u) \geq 0$, $G''_i(u) \leq 0$, $K''_i(u) \geq 0$. Using the generalized mean value theorem together with (A₂), (A₃) and (A₄), we obtain

$$(3.1) \quad f(t, u) \geq f(t, v) + F_u(t, v)(u - v) + \phi(t, v) - \phi(t, u),$$

$$(3.2) \quad g_i(u) \leq g_i(v) + G'_i(v)(u - v) + \psi_i(v) - \psi_i(u), \quad u, v \in \mathbb{R},$$

$$(3.3) \quad q_i(u) \geq q_i(v) + K'_i(v)(u - v) + \chi_i(v) - \chi_i(u), \quad u, v \in \mathbb{R}.$$

We set

$$\bar{F}(t, u, \alpha) = f(t, \alpha) + F_u(t\alpha)(u - \alpha) + \phi(t, \alpha) - \phi(t, u),$$

$$Q_i(u, \alpha) = q_i(\alpha) + K'_i(\alpha)(u - \alpha) + \chi_i(\alpha) - \chi_i(u),$$

$$\bar{g}_i(u(\gamma), \alpha, \beta) = g_i(\alpha(\gamma)) + G'_i(\beta(\gamma))(u(\gamma) - \alpha(\gamma)) + \psi_i(\alpha) - \psi_i(u),$$

and note that

$$\bar{F}_u(t, u, \alpha) < 0, \quad 0 \leq (\partial/\partial u)Q_i(u, \alpha) < 1, \quad 0 \leq (\partial/\partial u)\bar{g}_i(u(\gamma), \alpha, \beta) < 1.$$

Now, we fix $\alpha = \alpha_0$ and consider the problem

$$\begin{aligned} u''(t) + \sigma u'(t) + \bar{F}(t, u, \alpha_0) &= 0, & t \in [0, 1], \\ u(0) - \mu_1 u'(0) &= \bar{g}_1(u(\gamma), \alpha_0, \beta) + \int_0^{\gamma^-} Q_1(u(s), \alpha_0(s)) ds \\ &+ \int_{\gamma^+}^1 Q_1(u(s), \alpha_0(s)) ds, \\ u(1) + \mu_2 u'(1) &= \bar{g}_2(u(\gamma), \alpha_0, \beta) + \int_0^{\gamma^-} Q_2(u(s), \alpha_0(s)) ds \\ &+ \int_{\gamma^+}^1 Q_2(u(s), \alpha_0(s)) ds. \end{aligned} \tag{3.4}$$

As a first step, it will be shown that α_0, β are respectively lower and upper solutions of (3.4). Using (A₁) together with the fact that $\bar{F}(t, \alpha_0, \alpha_0) = f(t, \alpha_0)$, $\bar{g}_1(\alpha(\gamma), \alpha_0, \beta) = g_i(\alpha_0(\gamma))$ and $Q_i(\alpha_0, \alpha_0) = q_i(\alpha_0)$, we have

$$\alpha''_0(t) + \sigma \alpha'_0(t) + \bar{F}(t, \alpha_0, \alpha_0) = \alpha''_0(t) + \sigma \alpha'_0(t) + f(t, \alpha_0) \geq 0, \quad t \in [0, 1],$$

$$\begin{aligned} \alpha_0(0) - \mu_1 \alpha'_0(0) &\leq g_1(\alpha_0(\gamma)) + \int_0^{\gamma^-} q_1(\alpha_0(s)) ds + \int_{\gamma^+}^1 q_1(\alpha_0(s)) ds \\ &= \bar{g}_1(\alpha_0(\gamma), \alpha_0, \beta) + \int_0^{\gamma^-} Q_1(\alpha_0(s), \alpha_0(s)) ds \\ &+ \int_{\gamma^+}^1 Q_1(\alpha_0(s), \alpha_0(s)) ds, \end{aligned}$$

$$\begin{aligned} \alpha_0(1) + \mu_2 \alpha'_0(1) &\leq g_2(\alpha_0(\gamma)) + \int_0^{\gamma^-} q_2(\alpha_0(s)) ds + \int_{\gamma^+}^1 q_2(\alpha_0(s)) ds \\ &= \bar{g}_2(\alpha_0(\gamma), \alpha_0, \beta) + \int_0^{\gamma^-} Q_2(\alpha_0(s), \alpha_0(s)) ds \\ &+ \int_{\gamma^+}^1 Q_2(\alpha_0(s), \alpha_0(s)) ds \end{aligned}$$

and

$$\beta''(t) + \sigma \beta'(t) + \bar{F}(t, \beta, \alpha_0) \leq \beta''(t) + \sigma \beta'(t) + f(t, \beta) \leq 0, \quad t \in [0, 1].$$

Moreover, there exists $c_0, c_1 \in (\alpha_0(\gamma), \beta(\gamma))$ and $c_2, c_3 \in (\alpha_0, \beta)$ so that

$$\begin{aligned} &g_1(\beta(\gamma)) - \bar{g}_1(\beta(\gamma), \alpha_0, \beta) \\ &= g_1(\beta(\gamma)) - g_1(\alpha_0(\gamma)) - G'_1(\beta(\gamma))(\beta(\gamma) - \alpha_0(\gamma)) - \psi_1(\alpha_0(\gamma)) + \psi_1(\beta(\gamma)) \\ &= [g'_1(c_0) - g'_1(\beta(\gamma))](\beta(\gamma) - \alpha_0(\gamma)) + [\psi'_1(c_1) - \psi'_1(\beta(\gamma))](\beta(\gamma) - \alpha_0(\gamma)) \geq 0, \end{aligned}$$

$$\begin{aligned} &q_1(\beta(s)) - Q_1(\beta(s), \alpha_0(s)) \\ &= q_1(\beta(s)) - q_1(\alpha_0(s)) - K'_1(\alpha_0(s))(\beta(s) - \alpha_0(s)) - \chi_1(\alpha_0(s)) + \chi_1(\beta(s)) \\ &= [q'_1(c_2) - q'_1(\alpha_0(s))](\beta(s) - \alpha_0(s)) + [\chi'_1(c_3) - \chi'_1(\alpha_0(s))](\beta(s) - \alpha_0(s)) \geq 0 \end{aligned}$$

and consequently, we obtain

$$\begin{aligned} \beta(0) - \mu_1\beta'(0) &\geq g_1(\beta(\gamma)) + \int_0^{\gamma^-} q_1(\beta(s)) ds + \int_{\gamma^+}^1 q_1(\beta(s)) ds \\ &\geq \bar{g}_1(\beta(\gamma), \alpha_0, \beta) + \int_0^{\gamma^-} Q_1(\beta(s), \alpha_0(s)) ds + \int_{\gamma^+}^1 Q_1(\beta(s), \alpha_0(s)) ds. \end{aligned}$$

Similarly, it can be shown that

$$\beta(1) + \mu_2\beta'(1) \geq \bar{g}_2(\beta(\gamma), \alpha_0, \beta) + \int_0^{\gamma^-} Q_2(\beta(s), \alpha_0(s)) ds + \int_{\gamma^+}^1 Q_2(\beta(s), \alpha_0(s)) ds.$$

Thus we conclude that α_0 and β are respectively lower and upper solutions of (3.4). Hence, by Theorems 2.1 and 2.2, there exists the unique solution α_1 of (3.4) such that

$$\alpha_0(t) \leq \alpha_1(t) \leq \beta(t), \quad t \in [0, 1].$$

Next, we consider

$$\begin{aligned} &u''(t) + \sigma u'(t) + \bar{F}(t, u, \alpha_1) = 0, \quad t \in [0, 1], \\ &u(0) - \mu_1 u'(0) = \bar{g}_1(u(\gamma), \alpha_1, \beta) + \int_0^{\gamma^-} Q_1(u(s), \alpha_1(s)) ds \\ &\quad + \int_{\gamma^+}^1 Q_1(u(s), \alpha_1(s)) ds, \\ &u(1) + \mu_2 u'(1) = \bar{g}_2(u(\gamma), \alpha_1, \beta) + \int_0^{\gamma^-} Q_2(u(s), \alpha_1(s)) ds \\ &\quad + \int_{\gamma^+}^1 Q_2(u(s), \alpha_1(s)) ds. \end{aligned} \tag{3.5}$$

Using the earlier arguments, it can be shown that α_1 and β are lower and upper solutions of (3.5) respectively and by Theorems 2.1 and 2.2, there exists the unique solution α_2 of (3.5) such that

$$\alpha_1(t) \leq \alpha_2(t) \leq \beta(t), \quad t \in [0, 1].$$

Continuing this process successively yields a sequence $\{\alpha_n\}$ of solutions satisfying

$$\alpha_0(t) \leq \alpha_1(t) \leq \alpha_2(t) \leq \dots \leq \alpha_n(t) \leq \beta(t), \quad t \in [0, 1],$$

where the element α_n of the sequence $\{\alpha_n\}$ is a solution of the problem

$$\begin{aligned}
 u''(t) + \sigma u'(t) + \bar{F}(t, u, \alpha_{n-1}) &= 0, \quad t \in [0, 1], \\
 u(0) - \mu_1 u'(0) &= \bar{g}_1(u(\gamma), \alpha_{n-1}, \beta) + \int_0^{\gamma^-} Q_1(u(s), \alpha_{n-1}(s)) ds \\
 &\quad + \int_{\gamma^+}^1 Q_1(u(s), \alpha_{n-1}(s)) ds, \\
 u(1) + \mu_2 u'(1) &= \bar{g}_2(u(\gamma), \alpha_{n-1}, \beta) + \int_0^{\gamma^-} Q_2(u(s), \alpha_{n-1}(s)) ds \\
 &\quad + \int_{\gamma^+}^1 Q_2(u(s), \alpha_{n-1}(s)) ds
 \end{aligned}$$

and is given by

$$\begin{aligned}
 \alpha_n(t) &= \frac{-(1 - \sigma\mu_2)e^{-\sigma} + e^{-\sigma t}}{(1 + \sigma\mu_1) - (1 - \sigma\mu_2)e^{-\sigma}} \left[\bar{g}_1(\alpha_n(\gamma), \alpha_{n-1}, \beta) \right. \\
 &\quad \left. + \int_0^{\gamma^-} Q_1(\alpha_n(s), \alpha_{n-1}(s)) ds + \int_{\gamma^+}^1 Q_1(\alpha_n(s), \alpha_{n-1}(s)) ds \right] \\
 &\quad + \frac{(1 + \sigma\mu_1) - e^{-\sigma t}}{(1 + \sigma\mu_1) - (1 - \sigma\mu_2)e^{-\sigma}} \left[\bar{g}_2(\alpha_n(\gamma), \alpha_{n-1}, \beta) \right. \\
 &\quad \left. + \int_0^{\gamma^-} Q_2(\alpha_n(s), \alpha_{n-1}(s)) ds + \int_{\gamma^+}^1 Q_2(\alpha_n(s), \alpha_{n-1}(s)) ds \right] \\
 (3.6) \quad &+ \int_0^1 G(t, s) \bar{F}(s, \alpha_n(s), \alpha_{n-1}(s)) ds.
 \end{aligned}$$

Using the fact that $[0, 1]$ is compact and the monotone convergence of the sequence $\{\alpha_n\}$ is pointwise, it follows that the convergence of the sequence is uniform. If $u(t)$ is the limit point of the sequence, taking the limit $n \rightarrow \infty$ in (3.6), we obtain

$$\begin{aligned}
 u(t) &= \frac{-(1 - \sigma\mu_2)e^{-\sigma} + e^{-\sigma t}}{(1 + \sigma\mu_1) - (1 - \sigma\mu_2)e^{-\sigma}} \left[g_1(u(\gamma)) + \int_0^{\gamma^-} q_1(u(s)) ds + \int_{\gamma^+}^1 q_1(u(s)) ds \right] \\
 &\quad + \frac{(1 + \sigma\mu_1) - e^{-\sigma t}}{(1 + \sigma\mu_1) - (1 - \sigma\mu_2)e^{-\sigma}} \left[g_2(u(\gamma)) + \int_0^{\gamma^-} q_2(u(s)) ds + \int_{\gamma^+}^1 q_2(u(s)) ds \right] \\
 &\quad + \int_0^1 G(t, s) f(s, u(s)) ds.
 \end{aligned}$$

Thus, $u(t)$ is a solution of (2.1). Now, we show that the convergence of the sequence is quadratic. For that we set $\omega_n(t) = (u(t) - \alpha_n(t)) \geq 0$, $t \in [0, 1]$. In view of (A_2) ,

it follows by Taylor's theorem that

$$\begin{aligned}
 \omega_n''(t) + \sigma\omega_n'(t) &= u'' + \sigma u' - (\alpha_n'' + \sigma\alpha_n') = -f(t, u) + \bar{F}(t, \alpha_n, \alpha_{n-1}) \\
 &= -f(t, u) + f(t, \alpha_{n-1}) + F_u(t, \alpha_{n-1})(\alpha_n - \alpha_{n-1}) \\
 &\quad + \phi(t, \alpha_{n-1}) - \phi(t, \alpha_n) \\
 &= -f_u(t, c_4)(u - \alpha_{n-1}) - F_u(t, \alpha_{n-1})(u - \alpha_n) \\
 &\quad + F_u(t, \alpha_{n-1})(u - \alpha_{n-1}) - \phi_u(t, c_5)(\alpha_n - \alpha_{n-1}) \\
 &= [-f_u(t, c_4) + F_u(t, \alpha_{n-1}) - \phi_u(t, c_5)]\omega_{n-1} \\
 &\quad + [-F_u(t, \alpha_{n-1}) + \phi_u(t, c_5)]\omega_n \\
 &= [-F_u(t, c_4) + F_u(t, \alpha_{n-1}) + \phi_u(t, c_4) - \phi_u(t, c_5)]\omega_{n-1} \\
 &\quad + [-F_u(t, \alpha_{n-1}) + \phi_u(t, c_5)]\omega_n \\
 &\geq [-F_u(t, u) + F_u(t, \alpha_{n-1}) + \phi_u(t, \alpha_{n-1}) - \phi_u(t, \alpha_n)]\omega_{n-1} \\
 &\quad + [-F_u(t, \alpha_{n-1}) + \phi_u(t, \alpha_{n-1})]\omega_n \\
 (3.7) \quad &\geq [-F_{uu}(t, c_6) - \phi_{uu}(t, c_7)]\omega_{n-1}^2 - f_u(t, \alpha_{n-1})\omega_n \geq -A_1\|\omega_{n-1}\|^2,
 \end{aligned}$$

where $\alpha_{n-1} \leq c_4, c_6 \leq u, \alpha_{n-1} \leq c_5, c_7 \leq \alpha_n$, A is a bound on $\|F_{uu}\|$, B is a bound on $\|\phi_{uu}\|$ and $A_1 = A + B$. Further, we have

$$\begin{aligned}
 \omega_n(0) - \mu_1\omega_n'(0) &= g_1(u(\gamma)) - \bar{g}_1(\alpha_n(\gamma), \alpha_{n-1}, \beta) \\
 &\quad + \int_0^{\gamma^-} [q_1(u(s)) - Q_1(\alpha_n(s), \alpha_{n-1}(s))] ds \\
 &\quad + \int_{\gamma^+}^1 [q_1(u(s)) - Q_1(\alpha_n(s), \alpha_{n-1}(s))] ds \\
 &= g_1(u(\gamma)) - g_1(\alpha_{n-1}(\gamma)) - G_1'(\beta(\gamma))(\alpha_n - \alpha_{n-1}) \\
 &\quad + \int_0^{\gamma^-} [q_1(u(s)) - q_1(\alpha_{n-1}(s)) - K_1'(\alpha_{n-1}(s))(\alpha_n - \alpha_{n-1}) - \chi_1(\alpha_{n-1}) \\
 &\quad + \chi_1(\alpha_n)] ds + \int_{\gamma^+}^1 [q_1(u(s)) - q_1(\alpha_{n-1}(s)) - K_1'(\alpha_{n-1}(s))(\alpha_n - \alpha_{n-1}) \\
 &\quad - \chi_1(\alpha_{n-1}) + \chi_1(\alpha_n)] ds \\
 &\leq \left[\frac{1}{2}g_1''(\xi_1) + \psi_1''(\eta_2)\right]\omega_{n-1}^2(\gamma) + [G_1'(\beta(\gamma)) - \psi_1'(\eta_1)]\omega_n(\gamma) \\
 &\quad + \int_0^{\gamma^-} [(K_1'(\alpha_{n-1}(s)) - \chi_1'(\eta_3))\omega_n(s) + \left(\frac{1}{2}q_1''(\xi_2) + \chi_1''(\eta_4)\right)\omega_{n-1}^2(s)] ds \\
 &\quad + \int_{\gamma^+}^1 [(K_1'(\alpha_{n-1}(s)) - \chi_1'(\eta_3))\omega_n(s) + \left(\frac{1}{2}q_1''(\xi_2) + \chi_1''(\eta_4)\right)\omega_{n-1}^2(s)] ds
 \end{aligned}$$

and

$$\begin{aligned}
 \omega_n(1) + \mu_2 \omega'_n(1) &= g_2(u(\gamma)) - \bar{g}_2(\alpha_n(\gamma), \alpha_{n-1}, \beta) \\
 &+ \int_0^{\gamma^-} [q_2(u(s)) - Q_2(\alpha_n(s), \alpha_{n-1}(s))] ds \\
 &+ \int_{\gamma^+}^1 [q_2(u(s)) - Q_2(\alpha_n(s), \alpha_{n-1}(s))] ds \\
 &\leq \left[\frac{1}{2} g_2''(\xi_3) + \psi_2''(\eta_5) \right] \omega_{n-1}^2(\gamma) + [G_2'(\beta(\gamma)) - \psi_2'(\eta_6)] \omega_n(\gamma) \\
 &+ \int_0^{\gamma^-} \left[(K_2'(\alpha_{n-1}(s)) - \chi_2'(\eta_7)) \omega_n(s) + \left(\frac{1}{2} q_2''(\xi_4) + \chi_2''(\eta_8) \right) \omega_{n-1}^2(s) \right] ds \\
 &+ \int_{\gamma^+}^1 \left[(K_2'(\alpha_{n-1}(s)) - \chi_2'(\eta_7)) \omega_n(s) + \left(\frac{1}{2} q_2''(\xi_4) + \chi_2''(\eta_8) \right) \omega_{n-1}^2(s) \right] ds,
 \end{aligned}$$

where $\alpha_{n-1} \leq \xi_j \leq u, j = 1, \dots, 4, \alpha_{n-1} \leq \eta_\nu \leq \alpha_n \leq u, \nu = 1, \dots, 8$. In view of (A₃) and (A₄), there exists $\lambda_i < 1, \lambda_i^* < 1, M_i \geq 0$ and $M_i^* \geq 0$ such that $|G_i' - \psi_i'| \leq \lambda_i^*, |K_i' - \chi_i'| \leq \lambda_i, |\frac{1}{2} q_i'' + \chi_i''| \leq M_i$ and $|\frac{1}{2} g_i'' + \psi_i''| \leq M_i^*$. Letting $\lambda = \max\{\lambda_1, \lambda_2\}, \lambda^* = \max\{\lambda_1^*, \lambda_2^*\}, M^* = \max\{M_1^*, M_2^*\},$ and $M = \max\{M_1, M_2\},$ we get

$$(3.8) \quad \left\{ \begin{aligned}
 \omega_n(0) - \mu_1 \omega'_n(0) &\leq M^* \omega_{n-1}^2(\gamma) + \lambda^* \omega_n(\gamma) \\
 &+ \lambda \left[\int_0^{\gamma^-} \omega_n(s) ds + \int_{\gamma^+}^1 \omega_n(s) ds \right] \\
 &+ M \left[\int_0^{\gamma^-} \omega_{n-1}^2(s) ds + \int_{\gamma^+}^1 \omega_{n-1}^2(s) ds \right], \\
 \omega_n(1) + \mu_2 \omega'_n(1) &\leq M^* \omega_{n-1}^2(\gamma) + \lambda^* \omega_n(\gamma) \\
 &+ \lambda \left[\int_0^{\gamma^-} \omega_n(s) ds + \int_{\gamma^+}^1 \omega_n(s) ds \right] \\
 &+ M \left[\int_0^{\gamma^-} \omega_{n-1}^2(s) ds + \int_{\gamma^+}^1 \omega_{n-1}^2(s) ds \right].
 \end{aligned} \right.$$

Using the estimates (3.7) and (3.8), we obtain

$$\begin{aligned}
 \omega_n(t) &= \frac{-(1 - \sigma\mu_2)e^{-\sigma} + e^{-\sigma t}}{(1 + \sigma\mu_1) - (1 - \sigma\mu_2)e^{-\sigma}} \left(g_1(u(\gamma)) - \bar{g}_1(\alpha_n(\gamma), \alpha_{n-1}, \beta) \right. \\
 &+ \int_0^{\gamma^-} [q_1(u(s)) - Q_1(\alpha_n(s), \alpha_{n-1}(s))] ds \\
 &+ \left. \int_{\gamma^+}^1 [q_1(u(s)) - Q_1(\alpha_n(s), \alpha_{n-1}(s))] ds \right) \\
 &+ \frac{(1 + \sigma\mu_1) - e^{-\sigma t}}{(1 + \sigma\mu_1) - (1 - \sigma\mu_2)e^{-\sigma}} \left(g_2(u(\gamma)) - \bar{g}_2(\alpha_n(\gamma), \alpha_{n-1}, \beta) \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \int_0^{\gamma^-} [q_2(u(s)) - Q_2(\alpha_n(s), \alpha_{n-1}(s))] ds \\
 & + \int_{\gamma^+}^1 [q_2(u(s)) - Q_2(\alpha_n(s), \alpha_{n-1}(s))] ds \\
 & + \int_0^1 G(t, s) [f(s, u(s)) - \bar{F}(t, \alpha_n, \alpha_{n-1})] ds \\
 \leq & \frac{-(1 - \sigma\mu_2)e^{-\sigma} + e^{-\sigma t}}{(1 + \sigma\mu_1) - (1 - \sigma\mu_2)e^{-\sigma}} \left[M^* \omega_{n-1}^2(\gamma) + \lambda^* \omega_n(\gamma) \right. \\
 & + \lambda \left(\int_0^{\gamma^-} \omega_n(s) ds + \int_{\gamma^+}^1 \omega_n(s) ds \right) + M \left(\int_0^{\gamma^-} \omega_{n-1}^2(s) ds + \int_{\gamma^+}^1 \omega_{n-1}^2(s) ds \right) \Big] \\
 & + \frac{(1 + \sigma\mu_1) - e^{-\sigma t}}{(1 + \sigma\mu_1) - (1 - \sigma\mu_2)e^{-\sigma}} \left[M^* \omega_{n-1}^2(\gamma) + \lambda^* \omega_n(\gamma) \right. \\
 & + \lambda \left(\int_0^{\gamma^-} \omega_n(s) ds + \int_{\gamma^+}^1 \omega_n(s) ds \right) + M \left(\int_0^{\gamma^-} \omega_{n-1}^2(s) ds + \int_{\gamma^+}^1 \omega_{n-1}^2(s) ds \right) \Big] \\
 & - \int_0^1 G(t, s) [\omega_n''(s) + \sigma \omega_n'(s)] ds \\
 \leq & M^* \omega_{n-1}^2(\gamma) + \lambda^* \omega_n(\gamma) + \lambda \left(\int_0^{\gamma^-} \omega_n(s) ds + \int_{\gamma^+}^1 \omega_n(s) ds \right) \\
 & + M \left(\int_0^{\gamma^-} \omega_{n-1}^2(s) ds + \int_{\gamma^+}^1 \omega_{n-1}^2(s) ds \right) + A_1 \|\omega_{n-1}\|^2 \int_0^1 G(t, s) ds \\
 \leq & M^* \|\omega_{n-1}\|^2 + \lambda^* \|\omega_n\| + \lambda \|\omega_n\| + M \|\omega_{n-1}\|^2 + A_2 \|\omega_{n-1}\|^2 \\
 = & \lambda^{**} \|\omega_n\| + M^{**} \|\omega_{n-1}\|^2
 \end{aligned}$$

where A_2 provides a bound on $A_1 \int_0^1 G(t, s)$. We choose λ^* and λ so that $\lambda^{**} = \lambda^* + \lambda < 1$ and $M^{**} = M^* + M + A_2$. Taking the maximum over $[0, 1]$, we get

$$\|\omega_n\| \leq \frac{M^{**}}{1 - \lambda^{**}} \|\omega_{n-1}\|^2,$$

where $\|u\| = \max \{|u(t)| : t \in [0, 1]\}$. This establishes the quadratic convergence of the sequence of iterates. □

Remark. The results obtained in [2] appear as a special case of our results if we take $\gamma = 1/2$ in (2.1) and $\psi_i \equiv 0 \equiv \chi_i, i = 1, 2$ in the assumptions (A₃) and (A₄) of Theorem 3.1.

REFERENCES

[1] Ahmad, B., *A quasilinearization method for a class of integro-differential equations with mixed nonlinearities*, *Nonlinear Anal. Real World Appl.* **7** (2006), 997–1004.
 [2] Ahmad, B., Alsaedi, A., *Existence of approximate solutions of the forced Duffing equation with discontinuous type integral boundary conditions*, *Nonlinear Anal. Real World Appl.* **10** (2009), 358–367.

- [3] Ahmad, B., Alsaedi, A., Alghamdi, B., *Analytic approximation of solutions of the forced Duffing equation with integral boundary conditions*, *Nonlinear Anal. Real World Appl.* **9** (2008), 1727–1740.
- [4] Ahmad, B., Naz, U., Khan, R. A., *A higher order monotone iterative scheme for nonlinear Neumann boundary value problems*, *Bull. Korean Math. Soc.* **42** (2005), 17–22.
- [5] Ahmad, B., Nieto, J. J., *The monotone iterative technique for three-point second-order integrodifferential boundary value problems with p -Laplacian*, *Boundary Value Problems* **2007** (2007), 9pp., Article ID 57481, doi: 10.1155/2007/57481.
- [6] Ahmad, B., Nieto, J. J., *Existence and approximation of solutions for a class of nonlinear impulsive functional differential equations with anti-periodic boundary conditions*, *Nonlinear Anal.* **69** (2008), 3291–3298.
- [7] Ahmad, B., Nieto, J. J., Shahzad, N., *The Bellman-Kalaba-Lakshmikantham quasilinearization method for Neumann problems*, *J. Math. Anal. Appl.* **257** (2001), 356–363.
- [8] Ahmad, B., Sivasundaram, S., *The monotone iterative technique for impulsive hybrid set valued integro-differential equations*, *Nonlinear Anal.* **65** (2006), 2260–2276.
- [9] Bellman, R., Kalaba, R., *Quasilinearization and Nonlinear Boundary Value Problems*, Amer. Elsevier, New York, 1965.
- [10] Bouziani, A., Benouar, N. E., *Mixed problem with integral conditions for a third order parabolic equation*, *Kobe J. Math.* **15** (1998), 47–58.
- [11] Cabada, A., Nieto, J. J., *Rapid convergence of the iterative technique for first order initial value problems*, *Appl. Math. Comput.* **87** (1997), 217–226.
- [12] Cannon, J. R., *Encyclopedia of Math. and its Appl.*, ch. The one-dimensional heat equation, Addison-Wesley, Mento Park, CA, 1984.
- [13] Cannon, J. R., Esteva, S. Perez, Hoek, J. Van Der, *A Galerkin procedure for the diffusion equation subject to the specification of mass*, *SIAM. J. Numer. Anal.* **24** (1987), 499–515.
- [14] Choi, Y. S., Chan, K. Y., *A parabolic equation with nonlocal boundary conditions arising from electrochemistry*, *Nonlinear Anal.* **18** (1992), 317–331.
- [15] Denche, M., Marhoune, A. L., *Mixed problem with integral boundary condition for a high order mixed type partial differential equation*, *J. Appl. Math. Stoch. Anal.* **16** (2003), 69–79.
- [16] Ewing, R. E., Lin, T., *A class of parameter estimation techniques for fluid flow in porous media*, *Adv. Water Res.* **14** (1991), 89–97.
- [17] Formaggia, L., Nobile, F., Quarteroni, A., Veneziani, A., *Multiscale modelling of the circulatory system: a preliminary analysis*, *Comput. Vis. Sci.* **2** (1999), 75–83.
- [18] Ionkin, N. I., *Solution of a boundary value problem in heat condition with a nonclassical boundary condition*, *Differ. Uravn.* **13** (1977), 294–304.
- [19] Kartynnik, A. V., *Three-point boundary value problem with an integral space-variable condition for a second-order parabolic equation*, *Differential Equations* **26** (1990), 1160–1166.
- [20] Ladde, G. S., Lakshmikantham, V., Vatsala, A. S., *Monotone Iterative Techniques for Nonlinear Differential Equations*, Pitman, Boston, 1985.
- [21] Lakshmikantham, V., Nieto, J. J., *Generalized quasilinearization for nonlinear first order ordinary differential equations*, *Nonlinear Times & Digest* **2** (1995), 1–10.
- [22] Lakshmikantham, V., Vatsala, A. S., *Mathematics and its Applications, 440*, ch. Generalized Quasilinearization for Nonlinear Problems, Kluwer Academic Publishers, Dordrecht, 1998.
- [23] Nieto, J. J., Rodriguez-Lopez, R., *Monotone method for first-order functional differential equations*, *Comput. Math. Appl.* **52** (2006), 471–484.
- [24] Shi, P., *Weak solution to evolution problem with a nonlocal constraint*, *SIAM J. Math. Anal.* **24** (1993), 46–58.

- [25] Vatsala, A. S., Yang, J., *Monotone iterative technique for semilinear elliptic systems*, Boundary Value Probl. **2** (2005), 93–106.
- [26] Yurchuk, N. I., *Mixed problem with an integral condition for certain parabolic equations*, Differential Equations **22** (1986), 1457–1463.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE
KING ABDULAZIZ UNIVERSITY
P.O. BOX. 80257, JEDDAH 21589, SAUDI ARABIA
E-mail: aalsaedi@hotmail.com