






ISSN: 1715-9997

Canadian Journal of **pure & applied** **sciences** an international journal



SENRA
Academic Publishers
Burnaby, British Columbia

EDITOR

MZ Khan, SENRA Academic Publishers
Burnaby, British Columbia, Canada

ASSOCIATE EDITORS

Erol Hassan, University of Queensland
Gatton, Australia

Paul CH Li, Simon Fraser University
Burnaby, British Columbia, Canada

EDITORIAL STAFF

Jasen Nelson
Walter
Hao-Feng (howie) Lai

MANAGING DIRECTOR

Mak, SENRA Academic Publishers
Burnaby, British Columbia, Canada

The Canadian Journal of Pure and Applied Sciences (CJPAS-ISSN 1715-9997) is a peer reviewed multi-disciplinary specialist journal aimed at promoting research worldwide in Agricultural Sciences, Biological Sciences, Chemical Sciences, Computer and Mathematical Sciences, Engineering, Environmental Sciences, Medicine and Physics (all subjects).

Every effort is made by the editors, board of editorial advisors and publishers to see that no inaccurate or misleading data, opinions, or statements appear in this journal, they wish to make clear that data and opinions appearing in the articles are the sole responsibility of the contributor concerned. The CJPAS accept no responsibility for the misleading data, opinion or statements.

Editorial Office

E-mail: editor@cjpas.ca

SENRA Academic Publishers
7845 15th Street Burnaby
British Columbia V3N 3A3 Canada
www.cjpas.org
E-mail: senra@cjpas.ca

Volume 2, Number 1
Jan 2008

CANADIAN JOURNAL OF PURE AND APPLIED SCIENCES

Board of Editorial Advisors

- | | |
|---|---|
| Richard Callaghan
University of Calgary, AB, Canada | Sally Power
Imperial College London, UK |
| David T Cramb
University of Calgary, AB, Canada | Gordon McGregor Reid
North of England Zoological Society, UK |
| Matthew Cooper
Grand Valley State University, AWRI, Muskegon, MI, USA | Pratim K Chattaraj
Indian Institute of Technology, Kharagpur, India |
| Anatoly S Borisov
Kazan State University, Tatarstan, Russia | Andrew Alek Tuen
Institute of Biodiversity, Universiti Malaysia Sarawak, Malaysia |
| Ron Coley
Coley Water Resource & Environment Consultants, MB, Canada | Dale Wrubleski
Institute for Wetland and Waterfowl Research, Stonewall, MB, Canada |
| Chia-Chu Chiang
University of Arkansas at Little Rock, Arkansas, USA | Dietrich Schmidt-Vogt
Asian Institute of Technology, Thailand |
| Michael J Dreslik
Illinois Natural History, Champaign, IL, USA | Diganta Goswami
Indian Institute of Technology Guwahati, Assam, India |
| David Feder
University of Calgary, AB, Canada | M Iqbal Choudhary
HEJ Research Institute of Chemistry, Karachi, Pakistan |
| David M Gardiner
University of California, Irvine, CA, USA | Daniel Z Sui
Texas A&M University, TX, USA |
| Geoffrey J Hay
University of Calgary, AB, Canada | SS Alam
Indian Institute of Technology Kharagpur, India |
| Chen Haoan
Guangdong Institute for drug control, Guangzhou, China | Biagio Ricceri
University of Catania, Italy |
| Hiro Yoshi Ariga
Hokkaido University, Japan | Zhang Heming
Chemistry & Environment College, Normal University, China |
| Gongzhu Hu
Central Michigan University, Mount Pleasant, MI, USA | C Visvanathan
Asian Institute of Technology, Thailand |
| Moshe Inbar
University of Haifa at Qranim, Tivon, Israel | Indraneil Das
Universiti Malaysia, Sarawak, Malaysia |
| SA Isiorho
Indiana University - Purdue University, (IPFW), IN, USA | Gopal Das
Indian Institute of Technology, Guwahati, India |
| Bor-Luh Lin
University of Iowa, IA, USA | Melanie LJ Stiassny
American Museum of Natural History, New York, NY, USA |
| Jinfei Li
Guangdong Coastal Institute for Drug Control, Guangzhou, China | Kumlesh K Dev
Bio-Sciences Research Institute, University College Cork, Ireland. |
| Collen Kelly
Victoria University of Wellington, New Zealand | Borislava Gutarts
California State University, CA, USA |
| Hamid M.K.AL-Naimiy
University of Sharjah, UAE | Xiaobin Shen
University of Melbourne, Australia |
| AQ Khan
University of Illinois at Chicago, IL, USA | Maria V Kalevitch
Robert Morris University, PA, USA |
| Eric L Peters
Chicago State University, Chicago, IL, USA | Xing Jin
University of Hong Kong, Hong Kong |
| Roustam Latypov
Kazan State University, Kazan, Russia | Leszek Czuchajowski
University of Idaho, ID, USA |
| Frances CP Law
Simon Fraser University, Burnaby, BC, Canada | Basem S Attili
UAE University, UAE |
| Guangchun Lei
Ramsar Convention Secretariat, Switzerland | David K Chiu
University of Guelph, Ontario, Canada |
| Atif M Memon
University of Maryland, MD, USA | Gustavo Davico
University of Idaho, ID, USA |
| SR Nasyrov
Kazan State University, Kazan, Russia | Andrew V Sils
Georgia Southern University Statesboro, GA, USA |
| Russell A Nicholson
Simon Fraser University, Burnaby, BC, Canada | Charles S. Wong
University of Alberta, Canada |
| Shakeel A Khan
University of Karachi, Karachi, Pakistan | |

TIME DEPENDENT PROBLEM OF DIFFRACTION OF A SPHERICAL ACOUSTIC WAVE FROM AN ABSORBING HALF PLANE

*Bashir Ahmad, Ahmed Al-Saedi and Falleh Al-Solamy
 Department of Mathematics, Faculty of Science, King Abdulaziz University,
 P O Box 80257, Jeddah 21589, Saudi Arabia

ABSTRACT

The diffraction of a spherical acoustic wave from an absorbing half plane due to an arbitrary time dependent source distribution in the presence of a moving fluid is considered. The convolution integral appearing in the process of calculating inverse temporal transform has been evaluated asymptotically to present the diffracted field. This procedure is applicable to any type of time dependence provided the duration of incoming signal is small as compared to the emitted signal.

Keywords: Time dependent problem, spherical acoustic wave, absorbing half plane.

INTRODUCTION

The transient wave phenomenon is an important field of discussion in the wave motion theory which gives a more complete picture of the wave processes. The present ability to produce short electromagnetic pulses, which are being used as diagnostic tools for the study of implosion and other wave material applications, requires the development of new time dependent techniques. Stimulated by various applications that require the explicit treatment of time dependent effects, many scientists worked on time dependent problems, for instance, Friedlander (1958), Rienstra (1981), Jones (1986), Lakhtakia *et al.* (1987), Lakhtakia *et al.* (1989), Sun *et al.* (1991), Asghar *et al.* (1991), Asghar *et al.* (1998).

The objective of this paper is to discuss the general time dependent problem of diffraction of a spherical acoustic wave from an absorbing half plane in a moving fluid, which is perhaps a first attempt for this type of boundary. The temporal Fourier transform has been applied to obtain the transfer function in the transformed plane. Once the transfer function is available, an inverse temporal transform is used to obtain the results in time domain. The convolution technique is employed to calculate the inverse temporal transform and the resulting convolution integral has been evaluated approximately. This method of calculations can be used for various types of incoming signals whose amplitude characteristic is negligible in certain parts of the frequency axis (low pass filters). These filters are important in applications because the results obtained for these filters can be used to analyze more general filters, see Papoulis (1962). Finally, we employ this procedure to calculate the diffracted field due to an impulsive point source and triangular pulse from an absorbing half plane.

Formulation of the Problem

We consider the scattering of an acoustic wave from a semi-infinite absorbing plane occupying the position $y = 0, x \leq 0$. The plane is assumed to be of negligible thickness and satisfies the absorbing boundary condition $p - u_n z_a = 0$ on both sides of its surface, Morse and Ingard (1961). Here, p is the surface pressure and u_n the normal derivative of the perturbation velocity, z_a the acoustic impedance of the surface and n a normal pointing from the fluid into the surface. The whole system is assumed to be placed in a fluid moving with subsonic velocity U parallel to the $x -$ axis. The perturbation velocity \mathbf{u} of the irrotational sound wave can be written as $\mathbf{u} = \nabla \psi$. The resulting pressure in the sound field is given by

$$p = -\rho_0 \left(\frac{\partial}{\partial t} + U \frac{\partial}{\partial x} \right) \psi,$$

where ρ_0 is the density of the undisturbed stream. We consider a point source to be located at the position (x_0, y_0, z_0) and the time dependence is introduced through the function $f(t)$. Thus, we have to solve the following time dependent boundary value problem

$$\left[\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} - \left(\frac{1}{c} \frac{\partial}{\partial t} + M \frac{\partial}{\partial x} \right)^2 \right] \psi = \delta(x - x_0) \delta(y - y_0) \delta(z - z_0) f(t), \quad (1)$$

*Corresponding author email: bashir_qau@yahoo.com

$$\left\{ \frac{\partial}{\partial y} \mp \beta M \frac{\partial}{\partial x} \mp \frac{\beta}{c} \frac{\partial}{\partial t} \right\} \psi(x, 0^\pm, z; t) = 0, \quad (x < 0), \quad (2)$$

$$\frac{\partial}{\partial y} \psi(x, 0^+, z; t) = \frac{\partial}{\partial y} \psi(x, 0^-, z; t),$$

$$\psi(x, 0^+, z; t) = \psi(x, 0^-, z; t), \quad (x > 0), \quad (3)$$

where $\beta = \rho_0 c / z_a$, c is the velocity of sound and $M = U / c$ is the Mach number. For subsonic flow, $|M| < 1$ and $\text{Re } \beta > 0$.

Solution of the Problem

We define the temporal Fourier transform pair as

$$\begin{cases} \phi(x, y, z; w) = \int_{-\infty}^{\infty} \psi(x, y, z; t) e^{iwt} dt \\ \psi(x, y, z; t) = \frac{\text{Re}}{\pi} \int_0^{\infty} \phi(x, y, z; w) e^{-iwt} dw. \end{cases} \quad (4)$$

Now, taking temporal transform of Eqs.(1)-(3), we obtain

$$\left\{ (1 - M^2) \frac{\partial^2}{\partial x^2} + 2ikM \frac{\partial}{\partial x} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} + k^2 \right\} \phi = \delta(x - x_0) \delta(y - y_0) \delta(x - x_0) F(w), \quad (5)$$

$$\left\{ \frac{\partial}{\partial y} \mp \beta M \frac{\partial}{\partial x} \pm ik\beta \right\} \phi(x, 0^\pm, z; w) = 0, \quad (x < 0), \quad (6)$$

$$\begin{cases} \frac{\partial}{\partial y} \phi(x, 0^+, z; w) = \frac{\partial}{\partial y} \phi(x, 0^-, z; w), \\ \phi(x, 0^+, z; w) = \phi(x, 0^-, z; w), \end{cases} \quad (x > 0), \quad (7)$$

where $F(w)$ is temporal transform of $f(t)$ and its value is negligible outside an interval $(-\Omega, \Omega)$ (Papoulis, 1962) and $k = w / c$. We observe that the mathematical problem in the transformed w - plane is the same as in case of steady state (harmonic time dependent) problem, Asghar *et al.* (1991) except a multiplicative factor $F(w)$ in the right hand side of the wave equation. Thus, we can employ the results of Asghar *et al.* (1991) directly to write down the diffracted field as

$$f(x, y, z; w) = \frac{AF(w)}{\sqrt{w}} e^{-iw(MR_1 \cos \theta_1 - R_{12})/c_1 + i\pi/4}, \quad (8)$$

where

$$A = \sqrt{\frac{c_1}{(1 - M^2)}} \left\{ \frac{BR_{12} - BM \cos \theta_0 - 2 \sin \frac{\theta}{2} \sin \frac{\theta_0}{2}}{R + R_0} \right\} \times \frac{1}{4\pi \sqrt{2\pi R_0 R R_{12}} (\cos \theta + \cos \theta_0) L_+(K \xi_2 \cos \theta) L_-(-K \xi_2 \cos \theta_0)}, \quad (9)$$

$$X - X_0 = R_1 \cos \theta_1, c_1 = c \sqrt{(1 - M^2)}, k = K \sqrt{(1 - M^2)},$$

$$L(v) = 1 + \frac{B(K - Mv)}{\sqrt{K^2 \gamma^2 - v^2}} = L_-(v) L_+(v), \text{ see Asghar et al. (1991).}$$

Now, taking the inverse temporal transform of Eq.(8), we get

$$\psi(x, y, z; t) = A \frac{\text{Re}}{\pi} \int_0^{\infty} F(w) e^{-iw(MR_1 \cos \theta_1 - R_{12})/c_1 + i\pi/4} \frac{1}{\sqrt{w}} e^{-iwt} dw. \quad (10)$$

It is important to note that $L_+(Kx_2 \cos q)$ and $L_-(-Kx_2 \cos q)$ are independent of K . The integral appearing in Eq.(10) may be solved using the convolution theorem. Thus,

$$\psi(x, y, z; t) = \int_0^{\infty} f(t) g(t - \tau) d\tau, \quad (11)$$

where

$$g(t) = A \frac{\text{Re}}{\pi} \int_0^{\infty} e^{-iw(MR_1 \cos \theta_1 - R_{12})/c_1 + i\pi/4} \frac{1}{\sqrt{w}} e^{-iwt} dw = A \frac{\text{Re}}{\pi} \int_0^{\infty} G(w) e^{-iwt} dw = \frac{H\left(t - (R_{12} - MR_1 \cos \theta_1)/c_1\right)}{\sqrt{\pi\left(t - (R_{12} - MR_1 \cos \theta_1)/c_1\right)}}, \quad (12)$$

where $H(\cdot)$ is the Heaviside function. Now, we give a brief description of the approximate method for evaluating the convolution integral (11) between two functions $f(t)$ and $g(t)$. For that, Eq.(8) can be rewritten as

$$\phi(x, y, z; w) = F(w)G(w).$$

The p 'th moment of the function $f(t)$ denoted by m_p , is defined by

$$m_p = \int_{-\infty}^{\infty} t^p f(t) dt.$$

Using the moment theorem (Papoulis, 1962),

$$F(w) = \sum_{p=0}^{\infty} \frac{m_p}{p!} (-iw)^p.$$

Thus, the field $f(x, y, z; w)$ takes the form

$$f(x, y, z; w) = G(w) \sum_{p=0}^{\infty} \frac{m_p}{p!} (-iw)^p. \quad (13)$$

Since $(iw)^p G(w)$ is the Fourier transform of the p 'th derivative of $g(t)$, therefore the inverse temporal transform of Eq. (13) term wise yields

$$\begin{aligned} \psi(x, y, z; t) = m_0 g(t) - \frac{m_1}{1!} \frac{dg}{dt} + \dots \\ + (-1)^p \frac{m_p}{p!} \frac{d^p g}{dt^p}. \end{aligned} \quad (14)$$

Thus, $\psi(x, y, z; t)$ appears as a series in terms of derivatives of $g(t)$ and the moments of $f(t)$. For the validity of Eq. (14), it is assumed that Eq.(13) converges for every w . Even, if this happens, the convergence is slow and a large number of terms is required for a satisfactory evaluation of $f(t)$. However, if we assume that the temporal transform $G(w)$ is negligible outside an interval $(-\Omega, \Omega)$ and in this interval $F(w)$ is sufficiently smooth, so that for $|w| < W$, a small number of terms in Eq.(13) suffices to approximate adequately $F(w)$, that is,

$$F(w) \approx m_0 + \dots + \frac{m_k}{k!} (-iw)^k, \quad |w| < \Omega.$$

Thus, Eq.(14) takes the form

$$\begin{aligned} \psi(x, y, z; t) \approx m_0 g(t) - \frac{m_1}{1!} \frac{dg}{dt} + \dots \\ + (-1)^p \frac{m_p}{p!} \frac{d^p g}{dt^p}. \end{aligned} \quad (15)$$

The above assumption roughly stated in terms of time domain says that the duration of $f(t)$ should be small as compared to the duration of $g(t)$.

Examples

(1) Consider the impulsive point source $f(t) = d(t)$, where $d(t)$ is the Dirac delta function. In this case

$$m_0 = 1, \quad m_1 = m_2 = \dots = m_k = 0,$$

and the field $\psi(x, y, z; t)$ given by Eq. (15) takes the form

$$\psi(x, y, z; t) = \frac{H\left(t - (R_{12} - MR_1 \cos \theta_1)/c_1\right)}{\sqrt{\pi\left(t - (R_{12} - MR_1 \cos \theta_1)/c_1\right)}}. \quad (16)$$

Here, we observe that the diffracted field given by Eq. (16) corresponds to the field presented in Asghar *et al.* (1996), which proves the validity of this method.

(2) As a second example, we consider the triangular pulse

$$f(t) = \begin{cases} 1 - \frac{|t|}{T}, & \text{if } |t| < T, \\ 0, & \text{if } |t| > T, \end{cases}$$

whose temporal transform is

$$F(w) = \frac{4 \sin^2(Tw/2)}{Tw^2}.$$

The first three moments of $f(t)$ are given by

$$m_0 = T, \quad m_1 = 0, \quad m_2 = T^3/6.$$

Substituting these values in Eq. (15), we obtain the field

$$\psi(x, y, z; t) \approx \frac{AT}{\sqrt{\pi(t - R_{13})}} \left\{ 1 + \frac{T^2}{16(t - R_{13})^2} \right\},$$

where A is given by Eq. (9) and

$$R_{13} = (R_{12} - MR_1 \cos \theta_1)/c_1.$$

Concluding Remarks

In this paper, we have successfully employed the method of moments to present the asymptotic diffracted field from an absorbing half plane due to an arbitrary time dependent source with the assumption that the duration of incoming signal is small as compared to the emitted signal. As far as the validity of the method is concerned, we have provided a useful check in Example (1). It is interesting to note that the field due to a rigid half plane can be obtained by taking the absorption parameter b to be zero and the still fluid results by taking Mach number M equal to zero in Eq. (15) respectively.

REFERENCES

Asghar, S., Ayub, M., Ahmad, B. and Qadir, A. 1991. Diffraction of a cylindrical acoustic wave by an absorbing half plane in a moving fluid. *J. Acoust. Soc. America*. 89: 2080-2083.

Asghar, S., Hayat, T., Ayub, M. and Ahmad, B. 1998. Scattering of a spherical Gaussian pulse near an absorbing half plane. *Applied Acoustics*. 54: 323-338.

Asghar, S., Ahmad, B. and Ayub, M. 1991. Point source diffraction by an absorbing half plane. *IMA J. Appl. Math.* 46: 217-224.

Asghar, S., Ahmad, B. and Ayub, M. 1996. Diffraction from an absorbing half plane due to a finite cylindrical source. *Acustica-Acta Acustica* 82: 365-367.

Friedlander, FG. 1958. *Sound Pulses*. Cambridge University Press.

Jones, DS. 1986. *Acoustic and Electromagnetic Waves*, Clarendon Press, Oxford.

Lakhtakia, A., Varadan, VK. and Varadan, VV. 1987. Time-harmonic and time-dependent radiation by bifractal dipole arrays. *International Journal of Electronics*. 63: 819-824.

Lakhtakia, A., Varadan, VV. and Varadan, VK. 1989. Time-harmonic and time-dependent dyadic Green's functions for some uniaxial gyroelectromagnetic media. *Applied Optics*. 28: 1049-1052.

Morse, PM. and Ingard, KU. 1961. *Encyclopedia of Physics*, Springer-Verlag, Berlin, Vol. I (Acoustics).

Rienstra, SW. 1981. Sound diffraction at a trailing edge. *J. Fluid Mech.* 108: 443-460.

Papoulis, A. 1962. *The Fourier Integral and Its Applications*, Mcgraw-Hill Book Company, Inc. New

York, USA.

Sun, Z., Gimenez, G., Vray, D. and Denis, F. 1991. Calculation of the impulse response of a rigid sphere using the physical optic method and modal method jointly. *J. Acoust. Soc. America*. 89: 10-18.